
The genus *Oxydothis*: new palmicolous taxa and phylogenetic relationships within the *Xylariales*

Iman Hidayat^{1,2,*}, Rajesh Jeewon³, Chaiwat To-anun¹, and Kevin D. Hyde^{3*}

¹Department of Plant Pathology, Chiang Mai University, Chiang Mai, Thailand

²Mushroom Research Centre, Moo3 Ban Phadeng, Pa Pae, Chiang Mai, 50150 Thailand

³Centre for Research in Fungal Diversity, Department of Ecological & Biodiversity, The University of Hong Kong, Hong Kong SAR, PR China

Hidayat, I., Jeewon, R., To-anun, C. and Hyde, K.D. (2006). The genus *Oxydothis*: New palmicolous taxa and phylogenetic relationships within *Xylariales*. *Fungal Diversity* 23: 159-179.

Oxydothis (*Xylariales*) is an ascomycete genus, commonly encountered on decaying monocotyledons, such as palms. During our study on diversity of palmicolous fungi in northern Thailand, we encountered three new species of *Oxydothis*: *O. cyrtostachicola*, *O. inaequalis* and *O. wallichianensis*, and these are described and illustrated in this paper. The three novel species differ from other morphologically similar *Oxydothis* species in ascum shape and ostiole position, ascus ring, and ascospore morphology. Phylogenetic affiliations of the new taxa with members of related ascomycete families within the *Xylariales* are discussed based on morphology and nrDNA sequence data. The phylogenetic relationships of *Oxydothis* and its familial placement remain obscure based on the 28S nrDNA sequence analyses. Large sub unit nrDNA (28S) gene sequences do not provide significant phylogenetic information concerning the evolutionary relationships of these xylariaceous fungi. ITS nrDNA sequence analyses, however, indicate that *Oxydothis* is more closely related to members of the *Amphisphaeriaceae* than *Diatrypaceae* or *Xylariaceae*.

Key words: *Areaceae*, new species, palm fungi, ribosomal DNA, *Xylariales*

Introduction

Oxydothis Penz. & Sacc. (*Xylariales*) includes more than 70 species, which are saprobic, endophytic or parasitic on members of the *Gramineae*, *Liliaceae*, *Palmae* and *Pandanaceae* (Penzig and Saccardo, 1897; Hyde *et al.*, 2000). The genus is typified by *Oxydothis grisea* Penz. & Sacc. (Penzig and Saccardo, 1897) and delimited based on morphological features such as long

*Corresponding authors, Iman Hidayat, e-mail: hidayatiman@yahoo.com; Kevin D. Hyde, e-mail: kdhyde@hkucc.hku.hk

cylindrical asci with a J+ subapical apparatus; and long fusiform to filiform, hyaline, bicelled ascospores, which taper from the centre to spine-like ends, pointed or rounded processes (Penzig and Saccardo, 1897; Hyde, 1994). The familial placement of *Oxydothis*, however, is still tentative. There is uncertainty as to which morphological characters should be used to delimit the genus, as its taxonomic and phylogenetic relationships with other members of the *Xylariales* are still not resolved.

Based on morphological similarities with *Leiosphaerella* Höhn., and presence of horizontal and clypeate ascomata, *Oxydothis* was placed within the *Amphisphaeriaceae* (Muller and Arx, 1962; Muller and Arx, 1973; Wehmeyer 1976; Samuels and Rossman, 1987). The genus was, however, referred to the *Physosporaceae* (*Phyllachorales*) by Barr (1976) and later transferred to the *Hyponectriaceae* based on the perithecial ascomata with papillate ostiole, and cylindrical asci (Hawksworth *et al.*, 1995). A scanning Electron Microscopy (SEM) study of the ultrastructure of the *Oxydothis* ascus could not resolve the generic and familial affiliations of *Oxydothis* (Wong and Hyde, 1999). Wang and Hyde (1999) excluded *Oxydothis* from the *Hyponectriaceae* based on the arrangement of the ascomata, structure of the asci and ascospores, which are unlike *Hyponectria buxi*.

Phylogenetic analyses of 5.8S nrDNA and internal transcribed spacer (ITS2) sequence data have revealed that *Oxydothis* has evolutionary affiliations with members of the *Clypeosphaeriaceae* (Kang *et al.*, 1998). Although *Clypeosphaeriaceae* was found to be heterogeneous, Kang *et al.* (1999b, 2002) placed *Oxydothis* within the family. Phylogenies analyses of 28S and 18S nrDNA revealed closer affiliations of *Hyponectria* with *Oxydothis* and *Appendicospora*, although these relationships had poor statistical support (Smith *et al.*, 2003).

During our study on diversity of palmicolous fungi in northern Thailand, we collected four species of *Oxydothis*. There have been a number of novel ascomycetous taxa collected from Thailand (Chatmala *et al.*, 2004; Pinruan *et al.*, 2004; Somrithipol and Jones, 2005; Hunter *et al.*, 2006), but this is the first time we identified three new *Oxydothis* species. These are *Oxydothis cyrtostachicola* sp. nov. is described from decaying fronds of *Cyrtostachys renda* (*Arecaceae*), while *Oxydothis wallichianensis* sp. nov. and *O. inaequalis* sp. nov. are described from decaying leaves and fronds of *Wallichia siamensis* (*Arecaceae*), respectively. The new species are described and illustrated in this paper. In addition we performed phylogenetic analyses of partial 28S and ITS + 5.8S nrDNA sequence data to elucidate possible familial placement of *Oxydothis* within the *Xylariales*.

Materials and methods

Microscopic examination

Decaying fronds of *Wallichia siamensis* and *Cyrtostachys renda* were collected in northern Thailand during July and October 2005. The microscopic observation of *Oxydothis* spp. was performed as outlined in Shenoy *et al.* (2005) and Photita *et al.* (2005).

DNA extraction, amplification and sequencing

The total genomic DNA from the novel fungal species was extracted directly from the ascomata growing on the palm substrate, following a modified protocol of Hirata and Takamatsu (1996). Ascomata were picked up using fine forceps, and suspended in 500 µl of 5% Chelex solution (Bio-Rad, Richmond, Calif.). Ascomata were disrupted by means of a micropestle and content vortexed thoroughly for 1 min. Tube was incubated at a temperature of 100°C for 15 mins and vortexed again for 1 min to allow maximum disruption. The contents were centrifuged at a speed of 14000g for 30 seconds and the supernatant was transferred to a new tube. This sample was directly used as templates for PCR.

DNA amplification was performed by polymerase chain reaction (PCR). For partial 28S nrDNA amplification, LROR and LR5 primers (Vilgalys and Hester, 1990) were used, while ITS5 and ITS4 primers (White *et al.*, 1990) were used for ITS nrDNA amplification. The amplification procedure for ITS followed the protocol outlined in Jeewon *et al.* (2004) and Wang *et al.* (2005), while details for PCR conditions for 28S are outlined in Cai *et al.* (2005). Total reaction volume used was 25 µl. The PCR products spanning approximately 850 bp (partial 28S nrDNA) and 550 bp (ITS nrDNA), were checked on 1% agarose electrophoresis gels stained with ethidium bromide. The amplified PCR products were purified using minicolumns, purification resin and buffer according to the manufacturer's protocol (Amersham Biosciences, Catalog no. 27-9602-01). DNA sequencing was carried out using the above-mentioned primers in an Applied Biosystems 3730 DNA Analyzer at the Genome Research Centre, The University of Hong Kong.

Sequence alignment and phylogenetic analyses

For each species of *Oxydothis*, sequences obtained from the respective primers (LROR and LR5; ITS5 and ITS4) were aligned in Clustal X (Thomson **Table 1**. Sequences used in the analyses.

Species	Genbank accession no.	
	ITS nrDNA	28S nrDNA
<i>Oxydothis cyrtostachicola</i>	DQ 660334	DQ 660337
<i>Oxydothis daemonoropsicola</i>	DQ 660335	DQ 660338
<i>Oxydothis inaequalis</i>	DQ 660336	DQ 660339
<i>Amphisphaeria</i> sp.	AF375998	
<i>Arecophila bambusae</i>		AF452038
<i>Arecophila</i> sp.		AF452039
<i>Asordaria arctica</i>	AY681175	
<i>Bartalinia bischofiae</i>		AF382367
<i>Bartalinia laurina</i>	AF405302	AF382369
<i>Bartalinia robillardoides</i>	AF405301	
<i>Bertia moriformis</i>		AY695261
<i>Bionectria ochroleuca</i>		AY489716
<i>Bionectria pityrodes</i>		AY489728
<i>Cainia graminis</i>		AF431949
<i>Camarops tubulina</i>		AY346266
<i>Camarops ustulinooides</i>		AY346267
<i>Chaetomium globosum</i>		AF286403
<i>Clypeosphaeria mamillana</i>	AF009808	
<i>Cryptosphaeria pullmanensis</i>	AJ302419	
<i>Cryptosphaeria subcutanea</i>	AJ302420	
<i>Diatrype bullata</i>	DQ006946	
<i>Diatrype disciformis</i>		U47829
<i>Diatrype</i> sp.	DQ006957	
<i>Diatrype stigma</i>	AF192323	
<i>Discostroma fuscillum</i>	AF377284	
<i>Discostroma</i> sp.		AF382380
<i>Discostroma tosta</i>	AF009814	
<i>Discostroma tricellulare</i>	AF377285	
<i>Dothidea insculpta</i>		DQ247802
<i>Dothidea sambuci</i>		AY544681
<i>Eutypa armeniacae</i>	DQ006948	
<i>Eutypa lata</i>	DQ006944	
<i>Eutypa</i> sp.		AY346280
<i>Eutypella vitis</i>	DQ006943	
<i>Halorosellinia oceanica</i>		AY083822
<i>Kretzschmaria clavus</i>	AJ390434	
<i>Lepteutypa cupressi</i>		AF382379
<i>Monographella albescens</i>	AJ132506	

Table 1 continued. Sequences used in the analyses.

Species	Genbank accession no.	
	ITS nrDNA	28S nrDNA
<i>Monographella albescens</i>	AJ132509	
<i>Muscodor albus</i>	AY244622	
<i>Muscodor albus</i>	AY527044	
<i>Nitschkia grevillei</i>		AY346294
<i>Oxydothis frondicola</i>	AF009803	AY083835
<i>Pestalotiopsis versicolor</i>	DQ334862	AF382357
<i>Pestalotiopsis westerdijkii</i>	DQ137856	
<i>Seiridium cardinale</i>	AF377298	AF382377
<i>Seiridium unicorne</i>	AF377299	
<i>Sordaria fimicola</i>	AY681188	AY780079
<i>Truncatella angustata</i>	DQ093715	
<i>Ustulina deusta</i>	AF201718	
<i>Xylaria acuta</i>		AY544676
<i>Xylaria arbuscula</i>	AY183369	
<i>Xylaria hypoxylon</i>	AY327478	AY544648
<i>Xylaria mali</i>	AF163040	
<i>Xylaria polymorpha</i>	AF163042	
<i>Xylaria</i> sp.	AY315404	

et al., 1997) and Bioedit (Hall, 1999). In total, two datasets were analysed: Dataset based on 28 nrDNA sequences (Dataset I), and Dataset based on 5.8S nrDNA (Dataset II). Genbank accession numbers and taxa used are listed in Table 1.

Phylogenetic analyses were performed in PAUP* (Swofford, 2002). Ambiguously aligned sites were excluded from the all analyses. Unweighted parsimony (UP) and weighted parsimony (WP) analyses were performed. Gaps were treated as missing data. WP analyses were also performed using a symmetric step matrix generated with the program STMatrix version 2.2 (François Lutzoni and Stefan Zoller, Department of Biology, Duke University, Durham, NC).

Trees were inferred using the heuristic search option with 1000 random sequence additions. Maxtrees were unlimited, branches of zero length were collapsed and all multiple parsimonious trees were saved. Descriptive tree statistics (tree length [TL], consistency index [CI], retention index [RI], related consistency index [RC], homoplasy index [HI], and log likelihood [-ln L]) were calculated for trees generated under different optimality criteria. Clade stability was assessed in bootstrap analyses with 1000 replicates, each with 10 replicates of random stepwise addition of taxa. Random sequence addition was used in the bootstrap analyses. Kishino-Hasegawa tests (Kishino and Hasegawa, 1989) were performed in order to determine whether trees were

significantly different. Trees were figured in Treeview (Page 1996). Other details are outlined in Arenal *et al.* (2005) and Promputta *et al.* (2005).

The best-fit model of evolution was determined by MrModeltest2.2 (Posada and Crandall, 1998). Posterior probabilities (PP) (Rannala and Yang 1996; Zhaxybayeva and Gogarten 2002) were determined by Markov Chain Monte Carlo sampling (BMCMC) in MrBayes 3.0b4 (Huelsenbeck and Ronquist, 2001), using above estimated model of evolution. Six simultaneous Markov chains were run for 1,000,000 generations and trees were sampled every 100th generations (resulting 10,000 total trees). The first 1,000 trees that represented the burn-in phase of the analyses were discarded and the remaining 9,000 were used for calculating posterior probabilities (PP) in the majority rule consensus rule tree. The analyses were repeated five times starting from different random trees to ensure trees from the same tree space were being sampled during each analyses. Posterior probabilities equal to and above 95% were regarded as significant.

Results

Taxonomy

Oxydothis cyrtostachicola Hidayat, To-anun & K.D. Hyde, **sp. nov.**

Mycobank 510055

(Figs 1-4)

Etymology: In reference to the host genus, *Cyrtostachys*.

Ascomata 130–155 µm diam, 20–34 µm alta, immersa, subglobosa, ostiolata. Asci 102–120 × 12–13 µm, 8-spori, pedunculati, aparatu apicale J–, praediti Ascosporae 48–52 × 5–6 µm, hyalinae, fusiformis, bicellulares.


Ascomata forming under slightly raised, ellipsoidal regions on the host surface, black border, solitary or in groups 2–3; in section immersed, subglobose, ostiole eccentric, long axis horizontal to that of the host surface with neck at one end, forming *ca* 130–155 µm diam. × 20–34 µm high. *Peridium* comprised of 2–3 layers outer layers of oblong, dark-brown cells and with an additional inner layer of oblong, hyaline cells. *Paraphyses* deliquesce early. *Asci* 102–120 × 12–13 µm, 8-spored, unitunicate, clavate, short pedicellate, J–, refractive subapical ring, has a canal leading to the apex. *Ascospores* 48–52 × 5–6 µm, fusiform, 1-septate, hyaline, tapering gradually from the central septum to pointed processes, without spine-like form.

Anamorph: Unknown.

Habitat: Saprobic on *Cyrtostachys renda* fronds.

Known distribution: Northern Thailand.

Material examined: THAILAND: Chiang Mai, Chiang Mai University garden (latitude 18.84.00; longitude 98.97.00), on petioles of *Cyrtostachys renda*, 30 October 2005, FIH 151 (MRC 0007; **holotype** designated here).



Figs 1-4. Micrographs of *Oxydothis cyrtostachicola* (from holotype). **1.** Vertical section through the ascoma. **2.** Asci and ascular ring. **3-4.** Ascospores. Bars: 1 = 20 μm ; 3-4 = 5 μm .

Isotype: THAILAND: Chiang Mai, Chiang Mai University garden (latitude 18.84.00; longitude 98.97.00), on petioles of *Cyrtostachys renda*, October 30, 2005, FIH 151 (HKU (M) 17170).

***Oxydothis daemonoropsicola* J. Fröhl. & K.D. Hyde**


Material examined: THAILAND: Chiang Mai, Doi Suthep (latitude 18.48.00; longitude 98.56.00), decaying rachis of *Wallichia siamensis*, 21 July, 2005, Iman Hidayat FIH 019 (MRC 0005, 0006).

Oxydothis inaequalis* Hidayat, To-anun & K.D. Hyde, *sp. nov.

Mycobank 510054

(Figs 5-10)

Etymology: From the Latin *inaequalis* meaning “unequal, various” in reference to the ascospore processes that appear slightly unequal in length.



Figs 5-10. Micrographs of *Oxydothis inaequalis* (from holotype). **5.** Vertical section through the ascoma. **6.** Ascomata on the host surface. **7-8.** Ascospores. **9.** Ascus. **10.** Ascus ring. Bars: 5 = 110 μm ; 6 = 10 mm; 7 = 7 μm ; 8 = 7.5 μm ; 9 = 12 μm ; 10 = 4 μm .

Ascomata 110–120 µm diam, 22–30 µm alta, immersa, ellipsoida, ostiolata. *Asci* 200–285 × 11.25–13.75 µm, 8-spori, unitunicati, pedunculati, aparatu apicale J+, 3.5–4 µm alti, 3–3.5 µm diam. praediti *Ascosporae* 78–100 × 5–6.25 µm, 1-2 seiate, fusiformis, hyalinae, bicellulares.

Stromata 5–40 mm long × 5–10 mm wide, visible as blackened ellipsoidal regions on the host surface, lacking borders. *Ascomata* 110–120 µm diam. × 22–30 µm high, forming slightly raised domes, singly or clustered in groups up to 10 (mostly 2–3); in section immersed, ellipsoid, long axis horizontal to that of the host surface, papilla at one end curving upwards to the host surface. Stromatic tissue surrounds the ascomata within the host hypodermis. *Peridium* 10–12.5 µm thick, comprised of 2–3 layers; outer layers of oblong, dark-brown cells and sometimes with an additional inner layer of oblong, hyaline cells. *Paraphyses* deliquescent early, septate, ca 2.5 µm in diam. *Asci* 200–285 × 11.25–12.5 µm, 8-spored, unitunicate, cylindrical, short pedicellate, J+, 4–6(–7) µm high, 3–4 µm diam., wedge-shaped, subapical ring, apically truncate. *Ascospores* (75–) 78–87.5 (–100) × 5–7.5 µm, 1–2 seriate, fusiform, 1-septate, hyaline, tapering gradually to form long pointed processes. The ascospores processes are sometimes uneven which may make the septum appear slightly eccentric.

Anamorph: Unknown.

Habitat: Saprobic on the *Wallichia siamensis* fronds.

Known distribution: Northern Thailand.

Material examined: THAILAND, Chiang Mai, Doi Suthep (latitude 18.48.00; longitude 98.56.00), decaying rachis of *Wallichia siamensis*, 21 July 2005, Iman Hidayat FIH 018 (MRC 0004; **holotype** designated here),


Isotype: THAILAND, Chiang Mai, Doi Suthep (latitude 18.48.00; longitude 98.56.00), decaying rachis of *Wallichia siamensis*, 21 July 2005, Iman Hidayat FIH 018 (HKU (M) 17169).

Oxydothis wallichianensis Hidayat, To-anun & K.D. Hyde, **sp. nov.**
Mycobank No. 510053 (Figs 11-16)

Etymology: In reference to the host genus, *Wallichia*.

Ascomata 70–150 µm diam, 55–100 µm alta, subglobosa, ostiolata. *Asci* 87.5–125 × 10–15 µm, 8-spori, unitunicati, pedunculati, aparatu apicale J+, 2–2.5 µm alti, 3–3.5 µm diam. praediti. *Ascosporae* 32.5–55 × 6.25–7.5 µm, 1-2 seriate, fusiformis, hyaline, bicellulares.

Stromata 20–30 mm long × 15–25 mm wide, surrounded by ellipsoidal, brown borders. *Ascomata* 70–150 µm diam. × 55–100 µm high, stromata domes on the host surface, mostly clustered in groups of up to 18; in section immersed to erumpent, subglobose, papilla at one end curving upwards to the host surface. *Peridium* 5–8.75 µm thick comprised of 2–3 layers outer layers of oblong, dark-brown cells. *Asci* 87.5–125 × 10–15 µm, 8-spored, unitunicate, cylindrical, pedicellate, with a J+, 2–2.5 µm high × 3–3.5 µm diam., wedge-shaped, subapical ring. *Ascospores* 32.5–55 × 6.25–7.5 µm, 1–2-seriate,



Figs 11–16. Micrographs of *Oxydothis wallichianensis* (from the holotype). **11–12.** Ascomata on the host surface. **13.** Vertical section through the ascoma. **14.** Peridium. **15.** Ascospores. **16.** Asci and ascus ring. Bars: 11, 12 = 2 mm; 13, 16 = 50 μ m; 14 = 5 μ m; 15 = 10 μ m.

fusiform, 1-septate, hyaline, tapering abruptly near the ends to form long spine-like processes.

Anamorph: Unknown.

Habitat: Saprobic on *Wallichia siamensis* leaves.

Known distribution: Northern Thailand.

Material examined: THAILAND: Chiang Mai, Doi Suthep (latitude 18.48.00; longitude 98.56.00), on decaying leaves of *Wallichia siamensis*, 21 July 2005, Iman Hidayat FIH 010 (MRC 0002; **holotype** designated here).

Isotype: THAILAND: Chiang Mai, Doi Suthep (latitude 18.48.00; longitude 98.56.00), on decaying leaves of *Wallichia siamensis*, 21 July 2005, Iman Hidayat FIH 010 (HKU (M) 17174).

28S nrDNA phylogenies

The 28S nrDNA dataset consisted of 26 taxa. The other reference taxa included members of known ascomycete families of the *Sordariomycetes*. *Dothidea sambuci* and *D. insculpta* were the designated outgroups. The final dataset comprised 913 characters. The dataset was deposited in TreeBASE (<http://www.treebase.org/>) under the study accession no. S1603. Likelihood-ratio test in MrModeltest2.2 suggested that the best fit model of evolution for this dataset was SYM+I+G. Thirty-nine characters (ambiguous regions) were excluded in the analyses. In parsimony analyses, when gaps were treated as missing data, there were 505 constant characters, 95 parsimony-uninformative characters, and 274 parsimony-informative characters for both UP and WP. Two trees were obtained from UP and WP analyses. Based on K-H test ($P^* \geq 0.05$), these four trees were not significantly different (details not shown). The single parsimonious tree (TL = 899, CI = 0.600, RI = 0.688, RC = 0.412, HI = 0.400, $-\ln L = 5890.26251$) generated from WP analyses with *Dothidea sambuci* and *D. insculpta* as outgroups and treating gaps as missing data is shown in Fig. 17.

Bootstrap values (equal to or above 50%) based on 1000 replicates are shown on the upper branches, while values of the posterior probabilities (PP) resulted from BMBMC analyses are represented on the lower branches.

ITS nrDNA phylogenies

This dataset consisted of 37 taxa and analyses covered the ITS and 5.8S region. *Sordaria fimicola* and *Asordaria arctica* were the designated outgroups. The final dataset comprised 612 characters. The dataset was deposited in TreeBASE (<http://www.treebase.org/>) under the study accession no. S1602. Likelihood-ratio test in MrModeltest2.2 suggested that the best-fit model of evolution for this dataset was GTR+I+G. One hundred and six characters (ambiguous regions) were excluded in the analyses. In parsimony analyses, when gaps were treated as missing data, there were 199 constant characters, 36 parsimony-uninformative characters, and 271 parsimony-informative characters for both UP and WP. Four trees were generated from




Fig. 17. Phylogenetic tree based on partial 28S nrDNA sequence data showing placement of members of the *Oxydothis* within class *Sordariomycetes*. Bootstrap values $\geq 50\%$ from maximum parsimony analyses are shown above internodes and thickened branches indicate Bayesian analyses when posterior probabilities values $\geq 95\%$. Bootstrap values which are less than 50% shown by asterisk.




Fig. 18. Phylogenetic tree based on ITS nrDNA sequence data showing placement of members of the *Oxydothis* within the *Xylariales*. Bootstrap values $\geq 50\%$ from maximum parsimony analyses are shown above internodes and thickened branches indicate Bayesian analyses when posterior probabilities values $\geq 95\%$. Bootstrap values, which are less than 50% shown by an asterisk.

= 0.318, HI = 0.543, -ln L = 6699.36613) generated from WP analyses and treating gaps as missing data is shown in Fig. 18.

Bootstrap values (equal to or above 50%) based on 1000 replicates are shown on the upper branches, while values of the posterior probabilities (PP) resulted from BMBMC analyses are represented on the lower branches.

Discussion

Oxydothis is a common ascomycete genus, with 72 species which have been reported from monocotyledons and especially on palms (Hyde, 1994; Shenoy *et al.*, 2005). Fröhlich and Hyde (2000) reported that *Oxydothis* was the most common ascomycete genus occurring on palms. The basic taxonomic features of *Oxydothis* were discussed by Hyde (1994), who emphasized on several morphological features to delimit the species, i.e., ascoma orientation, ascal ring size and shape, and ascospore apex structure.

Table 2. Comparison of *Oxydothis wallichianensis* with similar *Oxydothis* species (measurements from Hyde, 1994c and Fröhlich and Hyde, 2000)

Taxa	Ascus		Ascal ring		Ascospores	
	Length (μm)	Width (μm)	Height (μm)	Width (μm)	Length (μm)	Width (μm)
<i>O. wallichianensis</i>	87.5–125	10–15	2–2.5	3–3.5	32.5–55	6.25–7.5
<i>O. batuapoensis</i>	83.5–117.5	(4.9–) 5.3–7.2	0.2–0.4	1.3–1.55 (–2)	33.9–48.8	2.9–3.4 (–4.1)
<i>O. dispariapicis</i>	224.2–277	14.1–17	2–3.4	2.4–3	(98–) 115–132.5	(6–) 7.3–8.8
<i>O. licualicola</i>	155–205	(10.5–) 12.5–15	1.75–2.5	3	60–80	5–7.5
<i>O. livistonicola</i>	<i>ca</i> 260	<i>ca</i> 10	1.2–1.6	2.4–3.2	74–96	6–8
<i>O. parvula</i>	110–130	8–10	1–1.6	2–2.8	49–62	4–6
<i>O. perangusta</i>	96–130	6.4–8	0.96–1.45	1.1–1.45	48–64	3–3.8
<i>O. elaeidis</i>	95–130	14–16	1.6	2.4–3.2	44–56	6–8
<i>O. sabalensis</i>	95–130	14–16	0.8–1	2.6–4	44–56	4–6

The three novel species differ from other morphologically similar *Oxydothis* species in ascomata shape and ostiole position, ascal ring, and ascospore morphology. The ascospores of *Oxydothis wallichianensis* are similar to those of eight other species and these taxa are compared in Table 2. *Oxydothis elaeidis* and *O. sabalensis* are the most similar species. The character that distinguishes *O. wallichianensis* from *O. elaeidis* is the ascal ring size, which is larger in the former ($2\text{--}2.5 \times 3\text{--}3.5 \mu\text{m}$ vs $1.6 \times 2.4\text{--}3.2 \mu\text{m}$). The grouping of ascomata (up to 20) under brown regions are also specific to

O. wallichianensis and distinguishes it from other similar *Oxydothis* species, such as *O. parvula*, *O. sabalensis*, *O. batuapoensis* where ascomata are mostly solitary.

Oxydothis asymmetrica is the only *Oxydothis* species with similar ascospores to those of *Oxydothis inaequalis*, both in size and shape. The significant characters that distinguish it from *O. inaequalis* are ascus size (200–285 × 11.25 μm vs. (6.8–) 8.1–11.9 (–12) μm) and ascus ring size (3.5–4 × 3–3.5 μm vs. 0.8–1.55 × 2.1–2.5 μm) (Table 3). *Oxydothis asymmetrica* also differs in having lenticular ascomata in section with a central pore, while *O. inaequalis* has ellipsoidal ascomata with an eccentric orientation.

Table 3. Comparison of *Oxydothis inaequalis* with *O. asymmetrica* (measurements from Hyde, 1994c; Fröhlich and Hyde, 2000)

Taxa	Ascus		Ascal ring		Ascospores	
	Length (μm)	Width (μm)	Height (μm)	Width (μm)	Length (μm)	Width (μm)
<i>O. inaequalis</i>	200-285	11.25–13.75	3.5-4	3–3.5	78-100	5-6.25
<i>O. asymmetrica</i>	157.1-208.8	(6.8-)8.1-11.9(-12)	0.8-1.55(-7)	2.1-2.5	67.5-81.5	4.7-.5(-7)

The third *Oxydothis* species from *Wallichia siamensis* is *Oxydothis daemonoropsicola*. The characteristics of ellipsoidal ascomata, long cylindrical asci, J+ ascus ring, and fusiform ascospores are typical of *O. daemonoropsicola*. The size of asci, ascus ring, and ascospores are also close to those of the type as reported by Fröhlich and Hyde (2000) (Table 4).

Table 4. Comparison of *Oxydothis* sp. FIH 019 with similar *Oxydothis* species (measurements from Hyde, 1994c; Fröhlich and Hyde, 2000).

Taxa	Ascus		Ascal ring		Ascospores	
	Length (μm)	Width (μm)	Height (μm)	Width (μm)	Length (μm)	Width (μm)
<i>Oxydothis</i> sp. FIH 019	225–255	12.5–13.75	2–2.25	2–3	95–105	5–6.25
<i>O. daemonoropsicola</i>	222.2–282.8	12.1–18.1	2.55–3.8	2.55–3.8	91.8–112.2	5.1–7.7
<i>O. megalospora</i>	224.5–292.8	11.7–16.6	3–7.5	3–5	103.7–126.9	5.6–7.6

Only four species of *Oxydothis* are characterized by a J– subapical ring: *O. ianei* (Taylor and Hyde, 2003), *O. livistonae*, *O. nonamyloidea* and *O. nontincta* (Fröhlich and Hyde, 2000). Species with J– subapical rings are distinguished by ascospore shape, size, and in having or lacking mucilage at

the ends. *Oxydothis cyrtostachicola* is characterized by fusiform ascospores, which taper gradually from the central septum to pointed processes, (not spine-like), and without mucilaginous drops at the ends. *O. cyrtostachicola* is closer to *O. ianei* and *O. nonamyloidea*, however, the asci and ascospores in *O. cyrtostachicola* are distinct from *O. ianei* and *O. nonamyloidea* in size. These taxa are compared in Table 5.

Table 5. Comparison of *Oxydothis cyrtostachicola* with J- *Oxydothis* species (measurements from Hyde, 1994e; Fröhlich and Hyde, 2000; Taylor and Hyde, 2003)

Taxa	Ascus		Ascospores	
	Length (µm)	Width (µm)	Length (µm)	Width (µm)
<i>O. cyrtostachicola</i>	102–120	12–13	48–52	5–6
<i>O. ianei</i>	108–174	7.2–9.6	56–68	2.6–3.8
<i>O. nonamyloidea</i>	205–260	18–22	94–115	3.5–4.5
<i>O. nontincta</i>	(136–)146–200	11–14	68–96	5–6.4(–7)
<i>O. livistonae</i>	Ca 300	11–14	150–170	4–5.5

Barr (1990) placed *Oxydothis* in the *Hyponectriaceae* based on its similarity with *Pemphidium*. *Oxydothis* is very similar to *Pemphidium* Mont. in ascomata, asci, and ascospore morphology. The latter, however, differs from *Oxydothis* in having a non-amyloid subapical ring and 1-celled, cylindrical ascospores with unusual polar appendages (Hyde *et al.*, 2000). Wehmeyer (1975) placed *Pemphidium* in the *Amphisphaeriaceae* and this placement was accepted by Eriksson and Hawksworth (1991), who emphasized more on ascus and paraphyses morphology for its inclusion in the family (non-amyloid apical ring in asci is not typical of the *Amphisphaeriaceae*). Wang and Hyde (1999) later excluded *Oxydothis* from the *Hyponectriaceae* based on arrangement of ascomata, structure of asci, and ascospores. They also excluded *Pemphidium* from the *Hyponectriaceae* due to few similarities with *Hyponectria*.

The non-amyloid subapical ring is not rare in *Oxydothis* since four reported *Oxydothis* species, *O. ianei* (Taylor and Hyde, 2003), *O. livistonae*, *O. nonamyloidea* and *O. nontincta* (Fröhlich and Hyde, 2000) have non-amyloid subapical rings.

Samuels and Rossman (1987) reported a *Selenosporella* anamorph for *Oxydothis selenosporellae*; however *Selenosporella* is also a synanamorph of *Iodosphaeria* (*Lasiosphaeriaceae*; Sordariales) (Samuels *et al.*, 1987). This may probably explain the distinctive evolutionary history of *Oxydothis* in relation to the members of the *Amphisphaeriaceae*, which are known to produce *Pestalotiopsis*-like anamorphs (Nag Raj, 1977; Kang *et al.*, 1999a;

Jeewon *et al.*, 2002; 2003). An ultrastructure of asci and ascospores of *Oxydothis* also suggested that *Oxydothis* is closer to the *Diatrypaceae* than the *Amphisphaeriaceae* (Wong and Hyde, 1999).

We used 28S nrDNA sequence data to confirm the placement of *Oxydothis* within the *Xylariales*. Molecular phylogenies in previous studies (Kang *et al.*, 1998, 1999b, 2002; Smith *et al.*, 2003) supported the placement of *Oxydothis* within the *Xylariales* (*Sordariomycetes*). Morphological characters such as presence of pseudostroma, ascoma perithecial, papillate ostiole, periphysate, cylindrical asci with J+ ring, and transversely septate ascospores also supports its placement within the *Xylariales*. Although *Oxydothis* species are nested in the *Xylariales* with 80% bootstrap support, there is insufficient evidence to place *Oxydothis* within any known families of the *Xylariales*. We note that molecular analyses does not support *Oxydothis* bears a monophyletic group and in addition *O. frondicola* did not cluster with other *Oxydothis* as would be expected based on morphological evidence.

Table 6. Morphological comparison of *Oxydothis* with *Amphisphaeriaceae* and *Hyponectriaceae*

Morphological characters	<i>Oxydothis</i>	<i>Amphisphaeriaceae</i>	<i>Hyponectriaceae</i>
Stromata	Pseudostroma	Crustose often clypeate	Reduced (clypeus absent)
Ascomata	Subglobose or often ellipsoidal and long axis parallel to the host surface	Globose	Subglobose
Asci shape	Long cylindrical	Cylindrical	Cylindric-clavate to clavate
Apical apparatus	J+ or J- subapical ring, variable in shapes (discoid, wedge shape, and cylindrical) and sizes	Usually J+, small	Small J+ or J-, apical ring
Ascospores colour	Hyaline	Hyaline to brown	Hyaline to pale brown
Ascospores shape	Long fusiform or filiform, gradually tapering from the centre to pointed processes, which may be spine-like, or with round ends	Ellipsoidal to fusiform	Various
Ascospores ornamentation	Often with small amounts of mucilage without germ pores	Rarely ornamented; however often with germ pores	Lacking ornaments

The association of *Oxydothis* with other known xylariaceous taxa receives very weak statistical support and most clades are not phylogenetically resolved. Therefore we refrain from making any systematic conclusions based on our 28S nrDNA sequence dataset. Another major observation in this study was the lack of statistical support for all the major nodes within the *Xylariales*. Even addition of more taxa with broader taxon sampling from all families failed to resolve some of the major clades. Similar results were obtained from previously published nrDNA phylogenies (Smith *et al.*, 2003; Duong, *et al.*, 2004; Bahl *et al.*, 2005). Parsimony analyses of the ITS dataset supported the monophyly of *Oxydothis* with 59% bootstrap support. *Oxydothis frondicola* and *O. daemonoropsicola* are closely related to each other, while *O. cyrtostachicola* and *O. inaequalis* share close phylogenetic affinities and both of these relationships are strongly supported. This is in accordance with their ascospore morphology. *Oxydothis frondicola* and *O. daemonoropsicola* produce filiform ascospores, while *O. cyrtostachicola* and *O. inaequalis* possess fusiform ascospores which tapering gradually from the central septum to pointed processes. Phylogenetic results also indicate that *Oxydothis* share close phylogenetic affinities to the *Amphisphaeriaceae* (74% bootstrap support). It is therefore highly plausible that *Oxydothis* is more closely related to members of the *Amphisphaeriaceae*, but whether it should be accommodated within this family as have been postulated by Muller and Arx, (1962; 1973), Wehmeyer (1976), Samuels and Rossman (1987), and Samuels *et al.*, (1987) based on morphology is still doubtful. On the other hand, whether *Oxydothis* should be accommodated in a new family will definitely need further detailed study with more species (including those from *Pemphidium*, *Leiosphaerella*, and *Iodosphaeria*) and sequence analyses based on protein-coding genes. Single gene phylogeny based on nrDNA is insufficient and inefficient in resolving phylogenetic relationships within the *Xylariales* (Smith *et al.*, 2003; Bahl *et al.*, 2005). We made several attempts to promote the formation of the anamorph of *Oxydothis* in culture and amplify partial RPB2 gene from a number of specimens but were unsuccessful.

Although the phylogeny of the *Hyponectriaceae* is still obscure, we propose that *Oxydothis* should be excluded from the *Hyponectriaceae* and *Amphisphaeriaceae*. Morphological characters such as ellipsoidal ascomata with long axis parallel to the host surface, long cylindrical asci with canal leads to the apex, variability of apical apparatus shape (discoid, wedge shape, and cylindrical) and size, ascospores with long fusiform or filiform with gradually tapering from the centre to pointed processes, support the exclusion from those families (Table 6). Furthermore, the discovery of *Oxydothis* anamorph will be the key factor in resolving its affinities.

Acknowledgments

The University of Hong Kong is thanked for providing DNA sequencing and other laboratory facilities to enable the molecular work. This research was funded by Hong Kong Research Grants Council (HKU 7322/04M) awarded to Dr. K.D. Hyde and Dr. R. Jeewon. Mushroom Research Centre (MRC) is acknowledged for providing Iman Hidayat with a scholarship to pursue his doctorate research at The University of Chiang Mai (Thailand). Belle Damodara Shenoy (University of Hong Kong) is thanked for his comments to improve the manuscript. Alvin Tang and Duong Minh Lam are thanked for laboratory assistance. Helen Leung and Heidi Kong are thanked for technical help. Jamjan Meeboon is thanked for collection assistance. Material was deposited in HKU(M) by agreement of MTA no. MPP 001.

References

- Arenal, F., Platas, G. and Peláez, F. (2005). *Preussia africana* and *Preussia isabelliae*, two new *Preussia* species based on morphological and molecular evidence. *Fungal Diversity* 20: 1-15.
- Bahl J., Jeewon, R. and Hyde, K.D. (2005). Phylogeny of *Rosellinia capetribulensis* sp. nov. and its allies (*Xylariaceae*). *Mycologia* 97: 1102-1110.
- Barr, M.E. (1978). *Buergenerula* and the *Physalosporaceae*. *Mycologia* 68: 611-621.
- Barr, M.E. (1990). Prodrum to nonlichenized, Pyrenomycetous members of Class Hymenoascomycetes. *Mycotaxon* 39: 43-104.
- Cai, L., Jeewon, R. and Hyde, K.D. (2005). Phylogenetic evaluation and taxonomic revision of *Schizothecium* based on ribosomal DNA and protein coding genes. *Fungal Diversity* 19: 1-17.
- Chatmala, I., Sakayaroj, J., Somrithipol, S. and Phongpaichit, S. (2004). Marine hyphomycetes of Thailand and *Cumulospora varia* sp. nov. *Fungal Diversity* 17: 1-9.
- Duong, L.M., Lumyong, S. Hyde, K.D. and Jeewon, R., (2004). *Emarcea castanopsicola* gen. et sp. nov. from Thailand, a new xylariaceous taxon based on morphology and DNA sequences. *Studies in Mycology* 50: 253-260.
- Eriksson, O. and Hawksworth, D.L. (1991). Outline to the ascomycetes – 1990. *Systema Ascomycetum* 9: 39-271.
- Fröhlich, J. and Hyde, K.D. (2000). *Palm Microfungi*. *Fungal Diversity Research Series*: 3: 1-393.
- Hall, T.A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analyses program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95-98.
- Hawksworth, D.L., Kirk, P.M., Sutton, B.C. and Pegler, D.N. (1995). *Ainsworth and Bisby's Dictionary of the Fungi*. CAB International, UK.
- Hirata, T. and Takamatsu, S. (1996). Nucleotides sequence diversity of rDNA internal transcribed spacers extracted from conidia and cleistothecia of several powdery mildew fungi. *Mycoscience* 37: 283-288.
- Huelsenbeck, J.P. and Ronquist (2001). MrBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754-755.
- Hunter, G.C., Crous, P.W., Wingfield, B.D., Pongpanich, K. and Wingfield, M.J. (2006). *Pseudocercospora flavomarginata* sp. nov. from *Eucalyptus* leaves in Thailand. *Fungal Diversity* 22: 71-90.
- Hyde, K.D. (1994b). Fungi from palms. VI. Reflection on *Oxydothis* and related genera. *Sydowia* 45: 204-225.

- Hyde, K.D. (1994e). Fungi from palms. XIII. The genus *Oxydothis*, a revision. *Sydowia* 46: 265-314.
- Jeewon, R., Liew, E.C.Y. and Hyde, K.D. (2002). Phylogenetic relationships of *Pestalotiopsis* and allied genera inferred from ribosomal DNA sequences and morphological characters. *Molecular Phylogenetics and Evolution* 25: 378-392.
- Jeewon, R., Liew, E.C.Y. and Hyde, K.D. (2003). Molecular systematics of the *Amphisphaeriaceae* based on cladistic analyses of partial LSU rDNA gene sequences. *Mycological Research* 107: 1392-1402.
- Jeewon, R., Liew, E.C.Y. and Hyde, K.D. (2004). Phylogenetic evaluation of species nomenclature of *Pestalotiopsis* in relation to host association. *Fungal Diversity* 17: 39-55.
- Kang, J.C., Kong, R.Y.C. and Hyde, K.D. (1998). Studies on the Amphisphaeriales: *Amphisphaeriaceae* (sensu stricto) and its phylogenetic relationships inferred from 5.8S rDNA and ITS2 sequences. *Fungal Diversity* 1: 147-157.
- Kang, J.C. Hyde, K.D. and Kong, R.Y.C. (1999a). Studies on Amphisphaeriales: the *Amphisphaeriaceae* (sensu stricto). *Mycological Research* 103: 53-64. .
- Kang, J.C. Hyde, K.D. and Kong, R.Y.C. (1999b). Studies on Amphisphaeriales 1: the *Clypeosphaeriaceae*. *Mycoscience* 40: 151-164.
- Kang, J.C. Hyde, K.D. and Kong, R.Y.C. (2002). *Phylogeny of Amphisphaeriaceae (sensu stricto)* and related taxa revisited based on nrDNA sequences. *Mycotaxon* 81: 321-330.
- Kishino, H. and Hasegawa, M. (1989). Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequence data, and the branching order in Hominidea. *Journal of Molecular Evolution* 29: 170-179.
- Kodsueb, R., Jeewon, R., Vijaykrishna, D., Hyde, K.D., McKenzie, E.H.C. and Lumyong, S. (2006). Systematic revision of *Tubeufiaceae* based on morphological and molecular data. *Fungal Diversity* 21: 104-129.
- Larget, B. and Simon, D. (1999). Markov chain Monte Carlo algorithms for the Bayesian analyses of phylogenetic trees. *Molecular Biology and Evolution* 16: 750-759.
- Muller, E. and von Arx, J.A. (1962). Die Gattungen der Didymospora Pyrenomycete. – *Beitrage der Kryptogamenflora Schweiz* 11: 1-992.
- Muller, E. and von Arx, J.A. (1976). Pyrenomycetes: Meliolales, Coronophorales, Sphaeriales. In: *The Fungi*. Vol. IVA (eds. G.C. Ainsworth, F.K. Sparrow and A.S. Sussman). Academic Press, New York: 87-132.
- Nag Raj, T.R. (1977). Miscellaneous microfungi. II. *Canadian Journal of Botany* 55: 757-765.
- Page, R.D.M. (1996). TreeView, an application to display phylogenetic trees on personal computers. *Computer Applications in the Biosciences* 12: 357-358.
- Penzig, O. and Saccardo, P.A. (1897). Diagnoses fungorum novorum in Insula Java collectorum. Series secunda. – *Malpighia* 11: 491-530.
- Photita, W., Taylor, P.W.J., Ford, R., Hyde, K.D. and Lumyong, S. (2005). Morphological and molecular characterization of *Colletotrichum* species from herbaceous plants in Thailand. *Fungal Diversity* 18: 117-133.
- Pinruan, U., McKenzie, E.H.C., Jones, E.B.G. and Hyde, K.D. (2004). Two new *Stachybotrys* species, and a key to the genus. *Fungal Diversity* 17: 145-157.
- Posada, D. and Crandall, K.A. (1998). Modeltest: testing the model of DNA substitution. *Bioinformatics* 14: 917-818.
- Promptutha, I., Jeewon, R., Lumyong, S., McKenzie, E.H.C. and Hyde, K.D. (2005). Ribosomal DNA fingerprinting in the identification of non sporulating endophytes from *Magnolia liliifera* (*Magnoliaceae*). *Fungal Diversity* 20: 167-186.

Fungal Diversity

- Rannala, B. and Yang, Z. (1996). Probability distribution of molecular evolutionary trees; a new method of phylogenetic inference. *Journal of Molecular Evolution* 43: 304-311.
- Ronquist, F. and Huelsenbeck, J.P. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572-1574.
- Samuels, G.J. and Rossman, A.Y. (1987). Studies in the *Amphisphaeriaceae* (*Sensu lato*) 2. *Leiosphaerella cocoës* and two new species of *Oxydothis* on palms. *Mycotaxon* 28: 461-471.
- Samuels, G.J., Müller, E. and Petrini, O. (1987). Studies in the *Amphisphaeriaceae* (*Sensu lato*) 3. new species of *Monographella* and *Pestalospaeria*, and two new genera. *Mycotaxon* 28: 473-499.
- Smith, G.J.D., Liew, E.C.Y. and Hyde, K.D. (2003). The Xylariales: a monophyletic order containing 7 families. *Fungal Diversity* 13: 175-208.
- Shenoy, B.D., Jeewon, R. and Hyde, K. D. (2005). *Oxydothis bambusicola*, a new ascomycete with a huge subapical ascus ring found on bamboo in Hong Kong. *Nova Hedwigia* 80: 511-518.
- Somrithipol, S. and Jones, E.B.G. (2005). An addition to the hyphomycete genus *Melanographium* from Thailand. *Fungal Diversity* 19: 137-144.
- Swofford, D.L. 2003. *PAUP*: Phylogenetic analyses using parsimony (* and other methods), version 4.0b 10*. Sinauer Associates, Sunderland, Massachusetts.
- Taylor, J.E. and Hyde, K.D. (2003). *Microfungi of tropical and temperate palms*. *Fungal Diversity Research Series*: 12: 1-458.
- Thomson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F. and Higgins, D.G. (1997). The Clustal_X windows interface: Flexible strategies for multiple sequences alignment aided by quality analyses tools. *Nucleic Acids Research* 25: 4876-4882.
- Vilgalys, R. and Hester, M. (1990). Rapid genetic identification and mapping enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238-4246.
- Wang, Y.Z. and Hyde, K.D. (1999). *Hyponectria buxi* with notes on the *Hyponectriaceae*. *Fungal Diversity* 3: 159-172.
- White, T.J., Burns, T., Lee, S. and Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: a Guide to Methods and Application* (eds. M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White (eds). Academic Press, San Diego: 315-322.
- Wehmeyer, L.E. (1975). The Pyrenomycetous fungi. *Mycologia Memoir* 6: 1-250.
- Wong, S.W. and Hyde, K.D. (1999). Ultrastructural observations on *Oxydothis alexandrarum*. *Fungal Diversity* 2: 181-188.
- Zhaxybayeva, O. and Gogarten, J.P. (2002). Bootstrap, Bayesian probability and maximum likelihood mapping exploring new tools for comparative genome analyses. *Genomics* 3: 1-15.

(Received 15 July 2006; Accepted 25 August 2006)